arXiv:cond-mat/9709182v2 12 Dec 1997

to be published *Modern Physics Letters B*

Polaron Dissociation at the Insulator-to-Metal Transition

P. Quémerais and S. Fratini
Laboratoire d'Etudes des Propriétés Electroniques des Solides,
(LEPES/CNRS), BP 166, 38042 Grenoble Cedex 9, France
(February 7, 2008)

Considering the long range Coulomb interactions between large polarons in dielectrics, we propose a model for their crystallization when no bipolarons are formed. As the density increases, the melting is examined at $T = OK$. One possibility is the delocalization towards a liquid state of polarons. However, we show that this cannot happen if the electron-phonon coupling is larger than some critical value. The other competing mechanism is the dissociation of the polarons themselves, favored owing to their large mass at strong coupling. Finally, we propose a phase diagram for the insulator-to metal transition as a function of the density and electron-phonon coupling.

Any theory of band structure for interacting fermions is based on the fact that the long range part of the Coulomb repulsion is screened and contained in collective modes, called plasmons [1]. However, for an idealized low density system the long range Coulomb repulsion dominates the kinetic energy, and Wigner [2] has shown that the carriers crystallize in that case. This does not occur for electrons in usual metals, owing to the high density of carriers.

For doped insulators, the carriers' density induced by doping can be extremely low. These materials can be classified into two types: polar and non-polar. Non-polar compounds are the usual covalent semiconductors which are easily doped up to the insulator-to-metal transition. Such materials are characterized by their dielectric constant κ, which does not depend on the frequency (the static dielectric constant ε_s and the high frequency dielectric constant ε_∞ are close together $\kappa \approx \varepsilon_s \approx \varepsilon_\infty$). The insulator-to-metal transition is due to the screening and ionization of the localized impurity states (the effects of disorder are neglected here). Mott [3] has given a criterium to evaluate the critical density above which the ground-state is metallic. Given the radius $R_{imp.} = \hbar^2\kappa/e^2m^*$ of the impurity bound state (m^* is the effective mass of the carrier), the system is metallic for doping densities n such that $n > n_c^M$, where $(n_c^M)^{1/3}R_{imp.} \approx 0.25$. Mott assumes that the long range Coulomb interactions between liberated carriers are also well screened as soon as $n > n_c$. This is correct because at these densities the kinetic energy of the carriers is much higher than the Coulomb energy $1/\kappa R_s$ (R_s is related to the density by $1/n = 4\pi R_s^3/3$), which is reduced by κ (which can be large), so that we are sure that the Wigner localization would occur only at densities much lower than the critical Mott density.

For polar insulators, such as some transition metal oxides, the situation is more complicated owing to the formation of polarons. When the static (ε_s) and high frequency (ε_∞) dielectric constants have an appreciable difference of magnitude, a free carrier is bound to the polarization that it creates around itself. The couple made of the carrier with its surrounding lattice polarization is called a polaron [4,5]. For the single polaron problem, one defines the dimensionless electron-phonon coupling $\alpha = \left(m^*/2\hbar^3\omega_{ir}\right)^{1/2} e^2/\tilde{\varepsilon}$. The constant $\tilde{\varepsilon}$, which is responsible for the polaron formation, is defined as $1/\tilde{\varepsilon} = 1/\varepsilon_\infty - 1/\varepsilon_s$, and ω_{ir} is the phonon frequency of the optical longitudinal mode. The polaron problem has been extensively addressed in many theroretical studies [6]. However, Feynman [7] has proposed a variational method using path integral calculations giving with a very good accuracy the ground state properties for any value of α. The idea is to use a simplified variational Hamiltonian to calculate quantities of interest:

$$H_0 = \frac{p^2}{2m^*} + \frac{P^2}{2M} + \frac{K}{2}(x - X)^2. \tag{1}$$

Here the electron with coordinate x and momentum p, is coupled to a fictitious particle with coordinate X and momentum P which simulates the polarization field. K and M are the variational parameters. As pointed out by Feynman, eq.(1) well imitates the polaron. In particular, M gives a good estimate to the true effective mass of the polaron : $M_P \approx m^* + M$, and the polaron radius can also be defined as $R_P \approx \sqrt{\hbar/2M_P(K/M_P)^{1/2}}$. Table 1 gives the rest mass M_P obtained by Feynman and Schultz [7] for different values of α. We notice that for large α, the polaron mass is much larger than the mass of the free carrier m^*.

If one considers now two polarons in the system, each electron (or hole) is attracted by the polarization induced by the other. On the other hand, there is also a direct repulsion between them. The resulting competition yields two different situations depending on the ratio $\eta = \varepsilon_\infty/\varepsilon_s$. Many theoretical calculations [8] have shown that if $\eta < \eta_c \approx 0.1$, a bound state between two polarons is formed, which is called a dielectric bipolaron. If this condition is not satisfied, the Coulomb repulsion dominates and both polarons repel as $1/\varepsilon_s d$, where d is their relative distance. We will assume from now on that we are in this last situation, which is often verified in real materials. All along this letter, we will consider materials which verify $0.1 < \eta < 0.5$.

On chemically doping at finite density polar dielectrics with $\eta > 0.1$, we are faced with three intricated problems: (i) the polaron formation, (ii) their binding to the impurity atoms (known to be the F-Center problem [9]), and (iii) the Coulomb repulsion between polarons if they are mobile. As already pointed out by one of us [10], the last point (iii) is essential to understand the physics of such systems. Since the polaron mass can be large (Table 1), the kinetic energy of a set of polarons which behaves as $1/M_P R_s^2$ can remain small compared to the Coulomb energy $1/\varepsilon_s R_s$ even at high density and high dielectric constant. To simplify the problem and focus mainly on the effects of the long range Coulomb interactions, we shall concentrate in this letter on an idealized system of polarons at finite density and interacting in a jellium of equal density (and opposite charge). At low density, they crystallize [10] as the electrons in the Wigner crystal. We will now address the problem of the melting of such a crystal as the density increases.

The strong electron-phonon coupling limit $\alpha \to \infty$, or equivalently the low phonon-frequency limit $\omega_{ir} \to 0$ can be first examined. It is well-known [4,9] that in this limit, the polaron is formed by an electron localized in a quasi-static polarization potential-well, which behaves as $-e^2/\tilde{\varepsilon} r$ at large distance, and is almost constant at short distance. The mass M_P of the polaron becomes infinite as ω_{ir} vanishes, while its energy E_P and radius R_P, which are finite, only depend on the dielectric constants and effective mass m^* of the electron. When the density increases, the polarons are unable to delocalize owing to their infinite mass. Thus, at some critical density, the electronic states localized within the polarization-wells must disappear for the same reason as the localized impurity states at the usual Mott transition: they are screened by liberated electrons. Making the same assumptions as Mott in the case of impurity states, we propose the following qualitative criterium for the dissociation of the polarons, valid for $\omega_{ir} \approx 0$, i.e $\alpha \to \infty$:

$$\left(n_c^P\right)^{1/3} \left(\frac{\tilde{\varepsilon}}{\varepsilon_\infty}\right) R_P \approx Const. \qquad (2)$$

The ratio $\tilde{\varepsilon}/\varepsilon_\infty$ takes into account the different dielectric constants appearing respectively in the calculation of the Thomas-Fermi length (ε_∞) and in the polaron radius ($\tilde{\varepsilon}$). The resulting state occuring for $n > n_c^P$ will be discussed later on. Numerically, on taking the values $\varepsilon_s = 30$, $\varepsilon_\infty = 5$, $m^* = m_e$ and $Const = 0.25$ [11], one obtains $n_c \approx 10^{19} cm^{-3}$.

The above consideration of the insulator-to-metal transition based on the polaron dissociation is *a priori* no more valid at finite electron-phonon coupling, since in that case the polarons could delocalize to form a liquid state instead of being dissociated. These two competing mechanisms for the insulator-to-metal transition are now more precisely examined at finite electron-phonon coupling and zero temperature. To modelize the situation at finite polaron density, we add to the host material a density n of electrons plus a *rigid* compensating jellium of same density. To account for the strong Coulomb interactions in a kind of mean-field approximation, starting from the low density limit, we take as a simplified model, a single electron localized within a uniformly positively charged and *polarizable* sphere of radius R_s, which is nothing but the Wigner crystal of polarons (see ref. [10] for details). As done by Wigner for the electron crystal, we neglect the dipole-dipole interactions between adjacent spheres. The hamiltonian describing such a system is then given by:

$$H = \frac{p_e^2}{2m^*} - \int\limits_{space} \mathbf{P}_{ir} \cdot \left(\mathbf{D}^- + \mathbf{D}^+\right) + \frac{1}{4\pi\varepsilon_\infty} \int\limits_{space} \mathbf{D}^- \cdot \mathbf{D}^+ + \frac{1}{8\pi\varepsilon_\infty} \int\limits_{space} \mathbf{D}^+ \cdot \mathbf{D}^+ + \frac{2\pi\tilde{\varepsilon}}{\omega_{ir}^2} \int\limits_{space} \left(\dot{P}_{ir}^2 + \omega_{ir}^2 P_{ir}^2\right), \qquad (3)$$

where $\mathbf{D}^+$ and $\mathbf{D}^-$ are the electric displacement due to the jellium and to electron respectively. $\mathbf{P}_{ir}$ is the polarization. We now introduce the polarization part which only responds to the electron motion : $\tilde{\mathbf{P}}_{ir} = \mathbf{P}_{ir} - (1/4\pi\tilde{\varepsilon})\mathbf{D}^+$ and $\dot{\tilde{\mathbf{P}}}_{ir} = \dot{\mathbf{P}}_{ir}$, and easily obtain:

$$H = C_0 + \frac{p^2}{2m^*} + \frac{1}{2} m^* \omega_W^2 x^2 - \int\limits_{space} \tilde{\mathbf{P}}_{ir} \cdot \mathbf{D}^- + \frac{2\pi\tilde{\varepsilon}}{\omega_{ir}^2} \int\limits_{space} \left(\dot{\tilde{P}}_{ir}^2 + \omega_{ir}^2 \tilde{P}_{ir}^2\right). \qquad (4)$$

$C_0 = -9e^2/10\varepsilon_s R_s$ is a constant part of the energy at fixed density, and $\omega_W^2 = e^2/m^* \varepsilon_s R_s^3$ is related to the usual plasmon frequency $\omega_p^2 = 4\pi n e^2/m^*$ by the relation $\omega_W^2 = \omega_p^2/3\varepsilon_s$. Equation (4) is nothing but the hamiltonian of a polaron, the electron of which is bound in a harmonic potential. Taking the limit $R_s \to \infty$, one recovers the usual single polaron hamiltonian, with $C_0 = \omega_W = 0$, $\mathbf{D}^+ = 0$, and $\tilde{\mathbf{P}}_{ir} = \mathbf{P}_{ir}$.

As previously proposed by Nozières *et al.* [12,13], the Lindemann criterium allows a good qualitative discussion for the melting of the electron Wigner crystal. The essential idea is that the crystallized state is unstable when the quantum fluctuations (zero point motion) of the electron positions exceed some critical value. More precisely, the crystal melts when $\left\langle \delta x^2 \right\rangle^{1/2} / R_s > \delta$, where δ is a phenomenological constant usually taken as $\delta \approx 1/4$. For the Wigner crystal, $\left\langle \delta x^2 \right\rangle = \hbar/2m^* \left(e^2/m^* R_s^3\right)^{1/2}$, and the critical value of R_s below which the crystal melts [14] is [12] : $R_c^W = 64 a_0$ (a_0 is the Bohr radius).

The application of this criterium to the polaron crystal is more complicated owing to the composite nature of the polaron. To be able to delocalize the polarons towards a liquid state, the quantum fluctuations for each single polaron moving as a whole, i.e. the motion of the electron together with the surrounding polarization, must become large in comparison with R_s. On the contrary, if the relative fluctuations of the electron with respect to the polarization become large, the polaron dissociation is favored. The Feynman model yields a natural way to evaluate the behavior of such quantities: since the polarization field is replaced by a particle with coordinate X, one easily calculates both the quantum fluctuations of the center of mass $R = (m^*x + MX)/(m^* + M)$ and the relative coordinate $r = x - X$. Thus, we have now two different Lindemann criteria:

(i) $\left\langle \delta R^2 \right\rangle^{1/2}/R_s > 1/4$ which allows the melting of the crystal towards a liquid state of polarons ;

(ii) $\left\langle \delta r^2 \right\rangle^{1/2}/R_s > 1/4$ for the polaron dissociation.

Both quantities are calculated by the same variational treatment, just taking instead of equation (2), the new Hamiltonian: $H_0^W = H_0 + \gamma x^2/2$, where γ, K and M are the new adjustable parameters. Once the lattice degrees of freedom are integrated out [7], the action for the model (4) is given by:

$$S = \frac{m^*}{2}\int dt\, \dot{x}^2(t) + \frac{m^*\omega_W^2}{2}\int dt\, x^2(t) - \frac{\alpha}{\sqrt{8}}\int\int dt\, ds\, \frac{e^{-|t-s|}}{|x(t) - x(s)|}, \tag{5}$$

where the integrals are calculated over a time interval T, $\hbar = 1$ and the frequency unit is ω_{ir}. The trial (quadratic) action corresponding to H_0^W is now

$$S_0^W = \frac{m^*}{2}\int dt\, \dot{x}^2(t) + \frac{\gamma}{2}\int dt\, x^2(t) + \frac{C}{2}\int\int dt\, ds\, [x(t) - x(s)]^2 e^{-w|t-s|}, \tag{6}$$

where $w^2 = K/M$ and $C = Kw/4$.

At low density, the two degrees of freedom r and R are uncoupled, and their frequencies are respectively ω_{int} and ω_{ext}. These are sketched in Fig. 1 as functions of n. The two corresponding quantum fluctuations are then given by $\delta r \approx (\hbar/2m^*\omega_{int})^{1/2}$ and $\delta R \approx (\hbar/2M_P\omega_{ext}))^{1/2}$. Fig.2.(a) shows the numerical values of the ratios (i) and (ii). One observes that for fixed α, while $\langle \delta r^2 \rangle^{1/2}/R_s$ increases with increasing density, $\langle \delta R^2 \rangle^{1/2}/R_s$ is bounded from above. In addition, its maximum value decreases with increasing α, so that above a certain critical value α^*, the value 1/4 is never reached and the crystal melting is prevented.

The situation encountered at infinite electron-phonon coupling (eq.(2)) is thus continued down to finite values of α. For $\alpha > \alpha^*$, *the liquid state of polarons cannot exist*. Physically, $\omega_{ext.}$ being limited by the frequency of the phonons themselves ($\omega_{ext.} \leq \omega_{ir}$, see Fig. 1), our result means that, when the electron-phonon coupling is too large, the polarization *cannot follow* the increase in kinetic energy induced by the doping through ω_W. The quantum fluctuations are then transferred to the *internal* degree of freedom. For that reason, the polarons dissociate at the transition.

Let us focus more precisely on the shape of $\langle \delta R^2 \rangle^{1/2}/R_s$ in Fig. 2(a). Starting at low density ($R_s \to \infty$, $\omega_W \approx 0$), the extra term $m^*\omega_W^2 x_e^2/2$ is a perturbation with respect to the single-polaron energy, and the polaron properties are almost unchanged as long as its radius is much smaller than the average distance between carriers. To a first approximation, we can then neglect the composite nature of the polaron and treat it as a rigid particle of mass M_P (the single polaron mass from Table 1) moving in a weak harmonic potential with a frequency $\omega_{ext} = \omega_W(m^*/M_P)^{1/2}$. The average quantum fluctuations in the jellium potential are given by the following expression

$$\langle \delta R^2 \rangle^{1/2} = \sqrt{\frac{\hbar}{2M_P(\omega_W\sqrt{m^*/M_P})}}, \tag{7}$$

which fits very well the low density part of $\langle \delta R^2 \rangle^{1/2}/R_s$ (right side of Fig. 2.(a)). This gives the critical $R_c^{(i)}$ corresponding to the first Lindemann criterium (i) in terms of $R_c^W = 64a_0$

$$R_c^{(i)} = R_c^W \frac{\epsilon_s}{(M_P/m^*)}. \tag{8}$$

If the mass of the carriers is strong enough, despite the static screening of the charges, the crystallisation is favored in comparison with the ordinary Wigner lattice. On the other hand, $\langle \delta r^2 \rangle^{1/2}$ in (ii) is nothing but the polaron radius, so that (ii) gives : $R_c^{(ii)} = 4R_P$.

At higher densities, the frequency of vibration of the polaron $\omega_{ext} = \omega_W(m^*/M_P)^{1/2}$ increases, due to the tightening of the jellium potential. Thus, if the density is such that $\omega_{ext} \approx \omega_{ir}$, i.e. $\omega_W \approx \omega_{ir}(M_P/m^*)^{1/2}$, the quenching of the polaron frequency is translated into an enhancement of the polaron mass, which can be described through a function $M_P(R_S)$. This is observed numerically, as shown in Fig. 2.(b) (the rest mass is recovered for $R_s \to \infty$). As a consequence, at high density, the quantum fluctuations of the polaron will be $(\hbar/2M_P(R_s)\omega_{ir})^{1/2}$ for the left part in Fig. 2.(a), while eq.(7) is valid for the right part. This gives rise to the bell shape of $\langle \delta R^2 \rangle^{1/2}/R_s$. This allows a simple evaluation of α^*: the maximum value of $\langle \delta R^2 \rangle^{1/2}/R_s$ will be obtained approximately by evaluating the low density formula eq.(4) at $\omega_W \approx \omega_{ir}(M_P/m^*)^{1/2}$. If we equate the result to the melting value 1/4 we get $(M_P/m^*)^{1/2}\alpha \approx 16(\epsilon_s/\tilde{\epsilon})$. As an example, taking $\epsilon_s = 30$

and $\epsilon_\infty = 5$, and using the values for M_P of Table 1, one obtains $\alpha^* \approx 9$. The variational result, without such approximations, gives $\alpha^* \simeq 7.5$.

The phase diagram Fig. 3 gives the calculated critical radii for crystal melting (i) and polaron dissociation (ii) as functions of the electron-phonon coupling α. Though the transition radii decrease linearly with ε_s and $\tilde{\varepsilon}$, the general shape of the diagram does not change in the studied range $0.1 < \varepsilon_\infty/\varepsilon_s < 0.5$, and so does the critical value α^* ($\alpha^* \approx 6-9$).

For small α the transition corresponds to the melting of the usual Wigner crystal of electrons weakly renormalized by the electron-phonon coupling. The corresponding metallic region ((I) in Fig.2) has been previously examined by Mahan [15]. Extension of his calculations in the metallic state have been performed by Devreese *et al.* at intermediate coupling, and more recently by Iadonisi *et al.* at strong coupling [15].

For intermediate coupling (say $5 < \alpha < \alpha^*$), it can be expected that the polarons remain well defined, yielding a liquid state of polarons below the transition (region (II) in Fig.2). The physics in that case may be well described by the Holstein model for non-interacting polarons [16,17].

For strong coupling ($\alpha > \alpha^*$), the crystal cannot melt towards a polaron liquid. At the transition, the polarons dissociate and the resulting metallic state (region (III) in Fig.2)) will present both strong electron-phonon and electron-electron interactions. A theory for such a system has been recently proposed by Castellani *et al.* [18]: it presents phase separation due to the electron-phonon coupling, which is prevented by the Coulomb repulsion. The resulting effects are strong charge density wave fluctuations, and a superconducting instability. Another possibility, already proposed by one of us [10] is that only part of the polarons dissociate to quasi-free electrons at the critical density. In that case, just above the insulator-to-metal transition the system would present a kind of electronic pseudo-separation between localized polarons and quasi-free electrons. However, at higher density, this pseudo-separation would disappear owing to screening effects, and in that case the system would be well represented by the theory of Castellani*et al.* [18]. This possibility will be examined in a further publication.

According to our theory, the possibility of polaron dissociation at the insulator-to-metal transition in real materials depends on the value of α, which is generally lower than $\alpha^* \approx 6-9$. However, any mechanism which increases the polaron mass M_P compared to the present 3D case, will shift α^* to lower values and make the dissociation of polarons physically reasonable. Moreover, the critical density for this dissociation will be increased compared to the present 3D case ($n_c^{3D} \approx 10^{19} cm^{-3}$, see eq. (2)). Two examples of such mechanisms can be mentioned: (a) a strong anisotropy of the effective mass m^*, which corresponds to considering two dimensional polarons; and (b) the formation of a polaron in a Mott-Hubbard insulator, which is associated to a spin-polaron [19], which mass increases exponentially with its size.

In conclusion, we have proposed a model which takes into account in a self-consistent way the effects of the Coulomb interaction for a system of polarons at finite densities. A new mechanism for the insulator-to-metal transition has been put in light, which corresponds to the polaron dissociation at strong electron-phonon coupling.

This work received financial support from the European Commission (contract no. ERBFMBICT 961230).

[1] D. Bohm and D. Pines, Phys. Rev., 85, 2, p.338 (1951)
[2] P. W. Wigner, Phys. Rev., 46, p.1002 (1934)
[3] N. F. Mott, Phil. Mag. 6, p. 2897 (1961)
[4] L. D. Landau, Phys. Z. Sowjetunion, 3, p.664 (1933)
[5] H. Frohlich, Adv. in Phys., 3, p.325 (1954)
[6] C. G. Kuper and G. D. Whitfield, *Polarons and excitons*, (Oliver and Boyd, Edinburgh and London, 1962)
[7] R. P. Feynman, Phys. Rev., 97, 3, p. 660, (1955) ; T. D. Schultz, Phys. Rev., 116, 3, p. 526, (1959)
[8] V. L. Vinetskii and M. S. Giterman, Soviet. Phys. JETP, 6, p.1796 (1958) ; G. Verbist, M.A. Smondyrev, F. M. Peeters, and J. T. Devreese, Phys. Rev. B, 45, 5262 (1992) ; F. Bassani, M. Geddo, G. Iadonisi and D. Ninno, Phys. Rev. B, 43, 5296 (1991)
[9] S. I. Pekar, AEC-tr-5575 Phys., State Publishing House of Technical and Theoritical Literature, Moscow (1961)
[10] P. Quémerais, Mod. Phys. Lett. B, 9, 25, p.1665 (1995)
[11] The constant *Const.* should be lower than the constant for the usual Mott criterium (≈ 0.25) owing to the difference of shape between a polarization potential well and an impurity potential well.
[12] P. Noziéres and D. Pines, Phys. Rev., 111, 2, P. 442 (1958)
[13] R.A. Coldwell-Horsfall, A.A. Maradudin, J.Math.Phys.1, 395 (1960)
[14] the value $R_c^W = 64$ overestimates the quantum fluctuations. See discussion in Ref. [12]
[15] G. Mahan, in *Polarons in Ionic Crystals and Polar semiconductors*, ed. J. Devreese (1972) ; L.F. Lemmens, J.T. Devreese and F. Brosens, Phys. Stat. Sol. 82, 439 (1977) ; G. Iadonisi, G. Capone, V. Cataudella and G. De Fillipis, Phys. Rev. B, 53, 13497 (1996)
[16] T. Holstein, Ann. of Phys., 8, 325 (1959)
[17] S. Ciuchi, F. De Pasquale, S. Fratini, D. Feinberg, Phys. Rev. B 56, 4494, (1997)
[18] C. Castellani, C. Di Castro and M. Grilli, Phys. Rev. Lett., 75, p.4650 (1995)
[19] A. S. Alexandrov and N. F. Mott, Rep. Prog. Phys. 57, 1197 (1994)

α	3	5	7	9	11
M_P/m^*	1.8	3.6	13.2	59.2	181

TABLE I. Polaron effective mass as given by Feynman.

FIG. 1. The two frequencies $\omega_{int.}$ and $\omega_{ext.}$ as functions of the density are represented in arbitrary units for $\alpha = 10$. The external frequency corresponding to the quantum fluctuations of the polaron as a whole saturates and is necessarily lower than ω_{ir}. The critical density n_c for the polaron dissociation (see text) is also quoted.

FIG. 2. (a) The ratios $\langle\delta R^2\rangle^{1/2}/R_s$ (bold line) and $\langle\delta r^2\rangle^{1/2}/R_s$ (dotted line) versus R_s/a_o at $\alpha = \alpha^*$ (see text), for $\epsilon_\infty = 5$, $\epsilon_s = 30$ and $m^* = m_e$. (b) The corresponding effective polaron mass M_P in units of m^*.

FIG. 3. Phase diagram for $\epsilon_\infty = 5$, $\epsilon_s = 30$ and $m^* = m_e$. Crystal melting (bold line) and polaron dissociation (broken line). The dotted line is the critical radius given by eq. (8). The arrow marks the result for the modified Mott criterium eq. (2). For regions I,II and III: see text.


ω_int
ω_ir
ω_ext
n_c
log(frequency)
log(density)


(a)
0.6
0.5
0.4
0.3
0.2
0.1
0
1
10
100
1000
10^4
$<\delta r^2>^{1/2}/R_s$
$<\delta R^2>^{1/2}/R_s$
R_s/a_0
(b)
10000
1000
100
10
M_P
1
10
100
1000
10^4
R_s/a_0



Wigner Crystal
Polaron Crystal
I
II
III
Metallic State
R_s/a_0
10^4
10^3
10^2
10^1
1
0
5
10
15
20
α